

Determination of Chemical Composition and Fumigant Insecticidal Activities of Essential Oils of Some Medicinal Plants Against the Adults of Cowpea Weevil, *Callosobruchus maculatus*

Kamuran KAYA[✉], Erdal SERTKAYA, İlhan ÜREMİŞ, Soner SOYLU

Department of Plant Protection, Mustafa Kemal University, Agricultural Faculty, Antakya, Hatay, Turkey
✉: soylu@mku.edu.tr

ABSTRACT

Chemical composition and fumigant insecticidal activities of essential oils obtained from oregano (*Origanum syriacum* L.), lavender (*Lavandula angustifolia* L.), sage (*Salvia officinalis* L.), fennel (*Foeniculum vulgare* Mill.) and laurel (*Laurus nobilis* L.) plants were investigated against the adults of cowpea weevil, *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae). Based on GC-MS analysis, total number of compounds determined in oregano, lavender, sage, fennel and laurel essential oils were 25, 33, 23, 23, 42, respectively. Carvacrol (66.66%), *trans*-anethol (79.17%), 1,8-cineole (50.0%), camphor (30.46%) and linalyl acetate (35.66%) were found as the most abundant compounds, respectively. Bioassay results revealed that essential oils, at 30.0 or 40.0 µg ml⁻¹ air, resulted in 100% adult mortality of *C. maculatus*. Insecticidal activity was increased in response to increased concentration of essential oil. Among the essential oils tested, oregano showed the highest fumigant toxicity at relatively lower concentration (30.0 µg ml⁻¹). Complete adult mortality (100%) caused by fennel, laurel, sage and lavender essential oils at the concentrations of 40.0 µg ml⁻¹ air. The LC₅₀ and LC₉₀ values for each essential oil were estimated by using probit analysis. The lowest LC₅₀ value was estimated for sage essential oil (8.79 µg ml⁻¹) followed by oregano (11.17 µg ml⁻¹), lavender (11.64 µg ml⁻¹), laurel (13.59 µg ml⁻¹) and fennel (17.46 µg ml⁻¹), respectively. The results revealed that plant essential oils might be used in research aiming the development of new environmental friendly control agents against stored-product pests such as cowpea weevil.

DOI:10.18016/ksudobil.386176

Article History

Received : 30.01.2018

Accepted : 06.04.2018

Keywords

Callosobruchus maculatus,
essential oil,
stored pests,
fumigant activity,
bioinsecticide

Research Article

Bazı Tıbbi Bitkilerden Elde Edilen Uçucu Yağların Kimyasal Bileşenlerinin Belirlenmesi ve Börülce Tohum Böceği, *Callosobruchus maculatus* Erginlerine Karşı Fümigant İnsektisidal Aktivitelerinin Belirlenmesi

ÖZET

Origanum (*Origanum syriacum* L.), lavanta (*Lavandula angustifolia* L.), adaçayı (*Salvia officinalis* L.), rezene (*Foeniculum vulgare* Mill.) ve defne (*Laurus nobilis* L.) bitkilerinden elde edilen uçucu yağların kimyasal bileşenleri ve fümigant insektisid etkinliği bakla tohum böceği *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae) erginlerine karşı araştırılmıştır. GC-MS analiz sonucuna göre, origanum, lavanta, adaçayı, rezene ve defne bitki uçucu yağlarında sırasıyla 25, 33, 23, 23, 42 adet bileşen belirlenmiştir. Carvacrol (%66.66), *trans*-anethol (%79.17), 1,8-cineole (%50.0), camphor (%30.46) ve linalyl acetate (%35.66) bu bitkilerin uçucu yağlarında ana bileşenler olarak belirlenmiştir. Biyolojik etkinlik çalışmaları tüm uçucu yağların 30.0 ve 40.0 µg ml⁻¹ hava konsantrasyonları *C. maculans* erginleri üzerinde %100 ölümlere neden olduğunu göstermiştir. Uçucu yağların insektisid etkinliği artan konsantrasyonlara bağlı olarak artış göstermiştir.

Makale Tarihiçesi

Geliş Tarihi : 30.01.2018

Kabul Tarihi : 06.04.2018

Anahtar Kelimeler

Callosobruchus maculatus,
uçucu yağ,
depo zararlısı,
fümigant etki,
biyoinsektisid

Araştırma Makalesi

Test edilen uçucu yağlar arasında, origanum uçucu yağı daha düşük konsantrasyonda ($30.0 \mu\text{g ml}^{-1}$ hava) en yüksek etkinlik göstermiştir. Rezene, defne, adaçayı ve lavanta uçucu yağları ise $40.0 \mu\text{gml}^{-1}$ hava konsantrasyonunda %100 ergin ölümlerine neden olmuştur. Her bir uçucu yağın LC_{50} ve LC_{90} değerleri probit analizi ile belirlenmiştir. En düşük LC_{50} değeri adaçayı ($8.79 \mu\text{g ml}^{-1}$) için belirlenmiş olup, bunu sırasıyla origanum ($11.17 \mu\text{g ml}^{-1}$), lavanta ($11.64 \mu\text{g ml}^{-1}$), defne ($13.59 \mu\text{g ml}^{-1}$) ve rezene ($17.46 \mu\text{g ml}^{-1}$) uçucu yağları takip etmiştir. Elde edilen sonuçlar bitki uçucu yağlarının börülce tohum böceği gibi depolanmış ürün zararlılarının kontrolünde yeni çevre dostu aktif maddelerin araştırıldığı çalışmalarda kullanılabileceğini göstermiştir.

To cite: Kaya K, Sertkaya E, Üremiş İ, Soylu S 2018. Determination of chemical composition and fumigant insecticidal activities of essential oils of some medicinal plants against the adults of cowpea weevil, *Callosobruchus maculatus*. KSÜ Tarım ve Doğa Derg 21(5): 708-714. DOI:10.18016/ksudobil.386176

INTRODUCTION

Cowpea (*Vigna unguiculata* (L.) Walp.), also called black-eyed pea, is important crop which is used as grain legume, vegetable or fodder crops. This crop is widely cultivated and consumed in warm regions around the world. Cowpea is an alternative source of dietary proteins and other essential nutrients (Brisibe et al., 2011). Production of cowpea is affected by pests both in the field and during storage. One of the most encountered stored pest in cowpea grain stack is the cowpea Bruchid *Callosobruchus maculatus* (F) (Coleoptera: Bruchidae). The insect infestation of cowpea seed during storage can cause economic loses and detrimental effect on seed germination of crop in several countries (Brisibe et al., 2011; Ileke and Olotuah, 2012). According to Caswell (1984) *C. maculatus* (F) populations persist and cause stock destruction as up to 50% over a period of eight months.

The most common management tactics applied to control for pests of stored grain are frequent use of synthetic chemical insecticides and fumigants. Effective and practical alternatives to chemical control and fumigants are currently not available. Besides, pesticide residues in grains, incidence of resistant insect populations and other negative effects of these chemicals to human health and environment, there is a public interest in development of alternative control methods such as development of bioactive chemical substances, conditioning of controlled and modified environment and integration of physical methods.

In this context, essential oils and their major chemical constituents have been explored by the scientific communities because of their excellent distribution (volatility), low toxicity in mammals and their rapid biodegradation capacity in ecosystem (Isman, 2011). Essential oils, are the volatile substance isolated by a physical process from odorous plant and categorized under plant secondary metabolites. Plant essential oils and major compounds were reported to have lethal effects, acting as pesticides against insects, mites and

various arthropods (Boulogne et al., 2012; El-Wakeil, 2013; Kedia et al., 2015). Essential oil and major compounds have also sublethal effects, acting as repellents and antifeedants in addition to their adverse effect on some biological parameters such as growth rate, life cycles and reproduction (Pascual-Villalobos 1996; Tunç and Erler, 2003; Erler and Tunç, 2005; Erler et al., 2006; Rattan, 2010). Since they can act as fumigant, their vapor action may also be very promising against pests of stored grain products because of their insecticidal properties (Rozman et al., 2007; Perez et al., 2010, Park et al., 2016).

This study was conducted to identify chemical constituents and to determine potential insecticidal effects of essential oils vapors of medicinal plants such as oregano (*Origanum syriacum* L.), sage (*Salvia officinalis* L.), lavender (*Lavandula angustifolia* L.), fennel (*Foeniculum vulgare* Mill.) and laurel (*Laurus nobilis* L.) against adult stage of cowpea weevil *C. maculatus*.

MATERIALS and METHODS

Insect material

C. maculatus adults were supplied from the naturally infested cowpea seeds in local warehouse in Hatay province. Healthy cowpea seeds were also purchased from the local market. The beetles were reared at Department of Plant Protection laboratory of Mustafa Kemal University (MKU) to obtain the parent stock. The insect culture was reared in the laboratory using clean and uninfested cowpea seeds which were free of insecticides. Clean and healthy cowpea seeds were disinfested by freezing at -20° for 48 h, with subsequent heating in an oven at 70°C for 24 h as described by Mkenda et al. (2015). The disinfested cowpea seeds were air dried at room temperature and stored in plastic box (25x16x11 cm, LxWxH).

Extraction of essential oils

The oregano, sage, lavender, fennel and laurel plants

were identified according to morphological characteristics and Flora of Turkey (Davis, 1988). Plant specimens were deposited in the herbarium collections of the Department of Plant Protection, MKU (No. OsKs7, FvN2, LnHb2, SoS21, LaSy5 and LaKp4). Naturally grown plants were collected from different districts of Hatay province (Samandağ, Defne and Antakya) situated in the Eastern Mediterranean region of Turkey. Approximately 200 g of milled air-dried leaves (for oregano, laurel, sage and lavender) or seeds (fennel) (150-200 g) together with 3 L double distilled water were placed into Clevenger apparatus (İldam, Ankara) for 3 h of steam-distillation. The resulting oils were separated over anhydrous sodium sulphate and stored in dark glass bottle and kept at 4 °C in the fridge for further use.

Analysis of chemical compositions of essential oils

The chemical compositions of the essential oils were determined using a Thermo Scientific Trace Ultra GC linked to a Thermo Scientific ISQ mass selective detector equipped with a TR-5 MS (Crosslinked 5% Phenyl Methyl Siloxane) capillary column (30 m x 0.25 mm i.d., 0.25 µm film thickness). The carrier gas was helium, at a ratio of 1.0 ml min⁻¹. The small volume of the samples (0.1 µl) was injected in split mode (50:1). The initial oven temperature was set to 50 °C, increased at a rate of 3 °C min⁻¹ to 210 °C and was maintained for 5 min. Identification of the spectra of major components of essential oils were determined by comparing their retention indices and mass spectra with authentic spectra present in the library (Wiley 275.L Registry of Mass Spectral Data) of the instrument (McLafferty and Staufner, 1989; Adams, 2007).

Determination of insecticidal activities of essential oils

The insecticidal activities of essential oils were determined by using transparent tubes (6 cm in height and 3.5 cm in diameters which offer 50 ml air space) as described previously (Sertkaya, 2013). Twenty cowpea weevil adults (2 to 4 days old) were introduced into each tube and 50 g cowpea seeds added and allowed weevils to settle for 30 min before adding the essential oils. Different concentrations of essential oils were applied on a filter paper strip (Whatman No.1) attached to under the surface of screw caps. Different concentrations of essential oil were prepared at the concentration of 4000.0 µg to 500.0 µg ml⁻¹ in DMSO and a 10-µl of each concentration was applied on filter paper strips (Whatman No.1) attached to under the surface of screw caps with a micropipette giving final concentrations of 5.0, 10.0, 15.0, 20.0, 30.0 and 40.0 µg ml⁻¹ air. Test tubes were, then, sealed with parafilm to prevent any loss of the essential oils. The control treatment included insects which were exposed to filter paper strip with 10 µl of DMSO. Three replicates of each treatments and control were arranged in

experiments.

The treated tubes were placed to incubator at 25 °C, 75 ±10%R.H., with a photoperiod of 12:12 (L:D) h (Shukla et al., 2011). Insect mortality was recorded under a stereo microscope daily after treatment. Insect mortalities were assessed for 120 min, as defined by lack of response to stroking with a paint-brush. Final mortality was calculated after 72 h by using Abbot's formula (Abbott, 1925).

Data analysis

The effect of different concentration of each essential oil on the adult mortality and the same concentration of different essential oils were analyzed with two-way analysis of variance (ANOVA). In all cases means were separated using Duncan's Multiple Range Test at $P \leq 0.05$. Probit analysis, in which probit-transformed percentage mortalities were regressed on log dose, was used to determine lethal concentrations (LC₅₀ and LC₉₀) values of each tested essential oil (Finney, 1971). All statistical analysis was performed using the SPSS statistical software program, version 17.0.

RESULTS and DISCUSSION

The chemical compositions of essential oils tested were presented in Table 1. Total number of component determined in oregano (*O. syriacum*), fennel (*F. vulgare*), laurel (*L. nobilis*), sage (*S. officinalis*) and lavender (*L. angustifolia*) was 25, 23, 42, 23 and 33, respectively. According GC-MS analysis, major constituent in the oregano essential oils was carvacrol (66.66%). The major component of *F. vulgare* was anethol (79.17%). The main component in *L. nobilis* oil was 1,8-cineole (50.0%). The major constituents in *S. officinalis* and *L. angustifolia* oils were camphor (30.46%) and linalyl acetate (35.66%), respectively.

The fumigant insecticidal activities of different concentrations of tested essential oils against *C. maculatus* adults were presented in Table 2. All essential oils resulted in 100% mortality of *C. maculatus* adult depending on the concentrations used. Among the tested essential oils, the highest fumigant insecticidal effect was shown by essential oil of oregano. Complete adult mortalities (100%) were observed at the concentration of 30.0 µg ml⁻¹. Essential oils of fennel, sage, laurel and lavender resulted in complete mortality at 40 µg ml⁻¹ concentration. In control treatments, adult mortality was not recorded. The lethal concentrations (LC₅₀ and LC₉₀) values (µg ml⁻¹) of each essential oil were determined by using probit analysis and given in Table 3. LC₅₀ values varied from 8.82 to 17.48 µg ml⁻¹ and LC₉₀ values ranged from 22.75 to 40.90 µg ml⁻¹ (Table 3). The lowest and highest LC₅₀ values were recorded for oils of sage (8.82 µg ml⁻¹) and fennel (17.48 µg ml⁻¹) essential oils, respectively. The lowest and highest LC₉₀ were recorded for oils of origanum (22.75 µg ml⁻¹) and fennel

(40.90 $\mu\text{g ml}^{-1}$) essential oils, respectively. The values determined for chi- (χ^2) values for all essential oils were found to be significant ($P > 0.15$). Since chemical pesticides cause detrimental effect on environment

and human being, several scientific researches concerning the use of environmentally friendly bio-insecticides, such as plant extracts, were initiated as a replacement to synthetic pesticides for integrated pest management (IPM) program (Bakkali et al., 2008).

Table 1. Major components detected in the essential oils used in this study

Essential oil	No. of components identified	Major components ^a identified
<i>Origanum syriacum</i> L.	25	carvacrol (66.66%), <i>p</i> -cymene (12.5%), γ -terpinene (12.4%)
<i>Foeniculum vulgare</i> Mill.	23	<i>trans</i> -anethole (79.17%), estragole (7.19%), limonene (6.61%)
<i>Laurus nobilis</i> L.	42	1,8-cineole (50.0%), α -terpineyl acetate (14.45%), sabinene (7.5%),
<i>Salvia officinalis</i> L.	23	Camphor (30.46%), Thujone (24.27%), 1,8 cineole (22.06%),
<i>Lavandula angustifolia</i> L.	33	linalyl acetate (35.66%), 1,8-cineole (32.65%), camphor (20.2%), lavandulyl acetate (6.19%),

^a Components showing a peak area of more than 5% relative to the total peak area on gas chromatography (GC) are listed in order of their highest relative peak area. Numbers are percentage of compound relative to total essential oil

Table 2. Percentage mortalities (%) and toxicity parameters of *Callosobruchus maculatus* after 24-h fumigation of tested essential oils at different concentrations

Concentrations ($\mu\text{g ml}^{-1}$ air)	Essential oils and % mortalities (\pm SE)				
	<i>O. syriacum</i>	<i>F. vulgare</i>	<i>S. officinalis</i>	<i>L. nobilis</i>	<i>L. angustifolia</i>
0.0	0.0 \pm 0.0a	0.0 \pm 0.0a	0.0 \pm 0.0a	0.0 \pm 0.0a	0.0 \pm 0.0a
5.00	13.33 \pm 3.3bBC	5.59 \pm 2.8aAB	29.35 \pm 1.0bD	3.03 \pm 3.0aA	20.71 \pm 3.3bCD
10.00	33.33 \pm 3.3cB	21.31 \pm 3.1bA	57.32 \pm 3.9cC	29.54 \pm 3.4bAB	31.21 \pm 2.7bAB
15.00	66.67 \pm 3.3dB	34.54 \pm 3.7cA	68.18 \pm 9.4cdB	57.93 \pm 7.0cB	62.43 \pm 5.3cB
20.00	83.33 \pm 3.3eA	55.81 \pm 1.2dA	78.18 \pm 13.1deA	75.46 \pm 7.7dA	74.85 \pm 10.5cdA
30.00	100.00 \pm 0.0fB	70.91 \pm 0.9eA	90.30 \pm 0.3efB	90.91 \pm 9.0deB	86.67 \pm 3.3deB
40.00	100.00 \pm 0.0f	100.00 \pm 0.0f	100.00 \pm 0.0f	100.00 \pm 0.0e	100.00 \pm 0.0e

Mean values (n=3 replicates with at least ten adult weevil per replicate) followed by the same small or capital letters within the column or row, respectively, are not significantly different according to Duncan's Multiple Range Test ($P \leq 0.05$).

Table 3. Lethal concentrations (LC) values of essential oils on *Callosobruchus maculatus* adult.

Probit Parameters	<i>O. syriacum</i>	<i>F. vulgare</i>	<i>S. officinalis</i>	<i>L. nobilis</i>	<i>L. angustifolia</i>
LC ₅₀ (CI 95%)	11.17 (9.56-12.78)	17.46 (15.13-20.16)	8.79 (6.71-10.68)	13.59 (11.85-15.36)	11.64 (9.63-13.69)
LC ₉₀ (CI 95%)	22.75 (19.26-28.97)	41.08 (33.13-57.38)	28.15 (22.06-41.68)	27.11 (23.11-34.26)	31.48 (25.19-44.35)
n	210	227	226	224	221
DF	4	4	4	4	4
Slope (\pm SE)	7.79 \pm 0.53	7.72 \pm 0.44	6.62 \pm 0.38	7.96 \pm 0.54	7.47 \pm 0.4
χ^2	4.274	6.436	2.421	1.057	4.858
P	0.370*	0.169*	0.659*	0.901*	0.302*
Y	-4.348+4.148x	-4.282+3.448x	-2.393+2.535x	-4.842+4.273x	-3.162+2.966x

LC concentration values are expressed as $\mu\text{g ml}^{-1}$ air

CI: confidence interval; n: number of adult insects used in the test; DF: degrees of freedom; SE: \pm standard error; χ^2 : chi-square; P: significance value; Y: Probit Equation, Probit (Pi): Intercept+BX (\log_{10} (dose));

*: significant ($P > 0.15$)

In previous studies, volatile components of essential oils were described to possess antimicrobial and pesticidal activities against different species of disease agent microorganisms and many pest arthropods

(Isman, 2000, Isman et al., 2011). Pyrethrum and neem oil which are the best established commercial natural insecticides were added the global pesticide market.

The insecticidal activities of essential oils of two *Cymbopogon* species on eggs and adults of two bruchids *C. maculatus* and *C. subinnotatus* exhibited a difference in bruchid susceptibility (Nyamador et al., 2010). Essential oil from *Cymbopogon giganteus* was reported to be more toxic against adult stages of these insect species. However, essential oil of *Cymbopogon nardus* showed the better ovicidal activity. Oviposition of females was strongly affected by the vapor of the essential oils used.

The chemical constituents and insecticidal activities of essential oils extracted from different plant species, such as eucalyptus, *Cymbopogon winterianus* and fennel plants were determined on adult population of *C. maculatus* by Gusmao et al. (2013). According to results obtained, all essential oils strongly displayed insecticidal effect on adults of *C. maculatus*, with varying LC₅₀ values according to essential oils used. Essential oils of *E. citriodora* and *C. winterianus* were found to be repellent while *F. vulgare* and *E. staigeriana* were classified as neutral against *C. maculatus* adults. According to results reported by Ebadollahi et al. (2012), the mean adult mortality rate of *C. maculatus* was significantly increased by increasing concentrations of essential oils of *Agastache foeniculum*, *A. graveolens*, *C. cyminum*, *F. vulgare* and *Satureja hortensis*. The chemical compositions of essential oils of *Eucalyptus globulus*, *E. radiata*, *Myrtus communis*, *Salvia officinalis*, *Laurus nobilis* and *Pistacia lentiscus* were investigated by GC/MS and their biological activity was assessed on *C. maculatus* on cowpea (Toudert-Taleb et al., 2014). All the oils tested by the scientists were found to be active and the contact dose of 12 µg l⁻¹ was found to be the most significant action on the inhibition of oviposition during the study. Panezail et al. (2015) recently described the insecticidal and repellent actions of medicinal plants including peppermint (*Mentha piperita*), sage (*Salvia officinalis*) and feverfew (*Tanacetum parthenium*), against two bruchid species *A. obtectus* and *C. maculatus*. They found that all of the tested plant essential oils had repellent and fatal effects against the adults of bruchid species. Among tested essential oils, maximum mortality for *A. obtectus* was recorded with the sage essential oil, whereas feverfew essential oils initiated the maximum mortality against *C. maculatus*. On the other hand, sage and feverfew exhibited maximum repellency against *C. maculatus* and *A. obtectus*, respectively. Similarly, Hashemi et al (2013) reported insecticidal activities of plant essential oil of sage (*Salvia leriifolia*) for *S. oryzae*, *C. maculatus* and *T. castaneum* adults. According to their result, adult mortalities of each pest species increased with the dose-depending manner and exposure time of essential oils.

To the best of our knowledge, this research is the first study regarding to *O. syriacum* and *L. angustifolia*

essential oils showing fumigant toxic activities on *C. maculatus* adults.

The number of components and their relative amount (%) present in essential oils show variation depending on plant species. Major components present in essential oils of different species of oregano, fennel, sage, laurel and lavender used in others study were also reported to contain carvacrol, anethole, camphor, and 1,8-cineole in previously published studies (Soylu et al., 2006; Sertkaya et al., 2010).

These major components have been described to have different insecticidal, acaricidal, herbicidal and antimicrobial activities against different species of insects, mites, weeds and plant disease agents (Isman, 2000; Bakkali et al., 2008; Park et al., 2016). Carvacrol, 1,8-cineole, camphor, anethole, terpinen and thymol, as major compounds of plant essential oils, were previously tested against different storage insects (Hamraoui and Regnault-Roger, 1997; Ahn et al., 1998; Kim and Ahn, 2001; Abeywickrama et al., 2006; Rozman et al., 2007; Suthisut et al., 2011, Park et al., 2016). According to their results, these individual compounds/constituents have lethal and sublethal deterrent activities on different species of stored grain pests. Monoterpenoids such as carvacrol, linalool and terpineol were stated to be more toxic in comparison to other compounds such as cymene, cinnamaldehyde, anethole against *A. obtectus* adults (Regnault-Roger and Hamraoui 1995). The results of Houghton et al. (2006) indicated that monoterpenoid compounds had an effect on insect mortality by re-regulating certain enzyme activity. In this frame, insecticidal effects of essential oils against adult cowpea bruchid mortalities might be the reason of the fumigant toxic effects of main constituents of essential oils.

It is well known from previously published studies that essential oils was reported to have antimicrobial, herbicidal and acaricidal efficiencies against fungal and bacterial disease agents, weeds and mites (Bakkali, 2008; Isman et al., 2011). Indeed, antimicrobial inhibitory activities of medicinal plants such as thyme, oregano, rosemary, lavender, fennel and laurel essential oils have been also described against important fungal and bacterial plant disease agents (Soylu et al., 2006, 2007; 2009; 2010).

In conclusion, essential oils obtained from naturally growing pharmaceutical plants in Hatay province revealed promising efficiencies for controlling *C. maculatus* adults. As a natural insecticides, plant essential oils possesses a wide range of desirable properties as alternative to synthetic pesticides because they are safer, have no toxicological effect against mammals and environment. However, possible use of these natural fumigants to control pest of stored-product needs further investigations to see whether these essential oils may be used in practice and also to assess the cost of their application in commercial way.

REFERENCES

- Abbott WS 1925. A method of computing the effectiveness of an insecticides. *J. Econ. Entomol.* 18: 265-267
- Abeywickrama K, Adhikari AACK, Paranagama P, Gamage CSP 2006. The efficacy of essential oil of *Alpinia calcarata* (Rosc.) and its major constituent, 1,8-cineole, as protectants of cowpea against *Callosobruchus maculatus* (F.) (Coleoptera : Bruchidae). *Can. J. Plant Sci.*, 86: 821-827.
- Adams RP 2007. Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy Allured Books. Carol Stream, IL. 698 pp.
- Ahn YJ, Lee SB, Lee HS, Kim GH 1998. Insecticidal and acaricidal activity of carvacrol and beta-thujaplicine derived from *Thujopsis dolabrata* var. *hondai* sawdust. *J. Chem. Ecol.*, 24: 81-90.
- Bakkali F, Averbeck S, Averbeck D, Waomar M 2008. Biological effects of essential oils—a review. *Food Chem. Toxicol.*, 46: 446–475.
- Boulogne I, Petit P, Ozier-Lafontaine H, Desfontaines L, Loranger-Merciris G 2012. Insecticidal and antifungal chemicals produced by plants: a review. *Environ. Chem. Lett.*, 10: 325-347.
- Brisibe EA, Adugbo SE, Ekanem U, Brisibe F, Figueira GM 2011. Controlling bruchid pests of stored cowpea seeds with dried leaves of *Artemisia annua* and two other common botanicals. *African J. Biotechnol.* 10: 9586–9592.
- Caswell GH 1984. The value of the pod in protecting cowpea seeds from attack by bruchid beetles. *Samaru J. Agric. Res.* 2: 49–55.
- Davis PH 1988. Flora of Turkey and the East Aegean Islands. Vol. I-X, Edinburgh Univ. Pres. (1965-1988).
- Ebadollahi A, Nouri-Ganbalani G, Hoseini SA, Sadeghi GR 2012. Insecticidal activity of essential oils of five aromatic plants against *Callosobruchus maculatus* F. (Coleoptera: Bruchidae) under laboratory conditions. *J. Essent. Oil Bearing Plants* 15: 256-262.
- El-Wakeil NE 2013. Botanical pesticides and their mode of action. *Gesunde Pflanzen* 65:125-149.
- Erler F, Uluğ I, Yalçinkaya B 2006. Repellent activity of five essential oils against *Culex pipiens*. *Fitoterapia*, 77: 491-494.
- Erler F, Tunç İ 2005. Monoterpenoids as fumigants against greenhouse pests: toxic, development and reproduction-inhibiting effects. *Z. Pflanzenk. Pflanzen.* (Journal of Plant Diseases and Protection), 112: 181-192.
- Finney DF 1971. Probit analysis, 3rd edn. University Press, Cambridge, p 333
- Gusmao NMS, de Oliveira JV, Navarro DMDF, Dutra KA, da Silva WA, Wanderley MJA 2013. Contact and fumigant toxicity and repellency of *Eucalyptus citriodora* Hook., *Eucalyptus staigeriana* F., *Cymbopogon winterianus* Jowitt and *Foeniculum vulgare* Mill. essential oils in the management of *Callosobruchus maculatus* (FABR.) (Coleoptera: Chrysomelidae, Bruchinae). *J. Stored Prod. Res.* 54: 41-47.
- Hamraoui A, Regnault-Roger C 1997. Monoterpenoids insecticidal effect comparison on two phytophagic insects, *Ceratitis capitata* and *Rhopalosiphum padi*. *Acta Botanica Gallica* 144: 413-417.
- Hashemi SM, Hosseini B, Estaji A, Hashemi SM, Hosseini B, Estaji A 2013. Chemical composition and insecticidal properties of the essential oil of *Salvia leriifolia* Benth (Lamiaceae) at two developmental stages. *J. Essent. Oil Bearing Plants*, 16: 806-816.
- Houghton PJ, Ren Y, Howes MJ 2006. Acetylcholinesterase inhibitors from plants and fungi. *Nat. Prod. Rep.* 23: 181–199.
- Ileke K, Olotuah O 2012. Bioactivity of *Anacardium occidentale* (L) and *Allium sativum* (L) powders and oils extracts against cowpea bruchid, *Callosobruchus maculatus* (Fab.) (Coleoptera: Chrysomelidae). *Int. J. Biol.* 4: 96–103.
- Isman MB 2000. Plant essential oils for pest and disease management. *Crop Protection* 19: 603–608.
- Isman MB, Miresmailli S, Machial C 2011. Commercial opportunities for pesticides based on plant essential oils in agriculture, industry and consumer products. *Phytochem. Rev.* 10: 197–204.
- Kedia A, Prakash B, Mishra PK, Singh P, Dubey NK 2015. Botanicals as eco-friendly biorational alternatives of synthetic pesticides against *Callosobruchus* spp. (Coleoptera: Bruchidae)-a review. *J. Food Sci. Technol-Mysore* 52: 1239-1257.
- Kim DH, Ahn YJ 2001. Contact and fumigant activities of constituents of *Foeniculum vulgare* fruit against three coleopteran stored-product insects. *Pest Manag. Sci.* 57: 301-306.
- McLafferty FW, Staufner DB 1989. The Willey N.B.S. Registry of mass spectral data. New York: John Wiley and Sons.
- Mkenda PA, Stevenson PC, Ndakidemi P, Farman DI, Belmain SR 2015. Contact and fumigant toxicity of five pesticidal plants against *Callosobruchus maculatus* (Coleoptera: Chrysomelidae) in stored cowpea (*Vigna unguiculata*). *Int. J. Tropical Insect Sci.* 35: 172–184.
- Nyamador WS, Ketoh GK, Amevoin K, Nuto Y, Koumaglo HK, Glitho IA 2010. Variation in the susceptibility of two *Callosobruchus* species to essential oils. *J. Stored Prod. Res.* 46: 48–51
- Panezail GM, Jabeen, R, Khetran M, Ijaz A, Rafeeq M, Bukhari FA, Awan MA, Hameed T, Tariq MM 2015. Insecticidal action of three plants extracts against cowpea weevil, *Callosobruchus maculatus* (F) and bean weevil, *Acanthoscelides obtectus* Say. *Pak. J. Zool.* 47: 899-902.

- Park CG, Shin E, Kim J 2016. Insecticidal activities of essential oils, *Gaultheria fragrantissima* and *Illicium verum*, their components and analogs against *Callosobruchus chinensis* adults. *J. Asia-Pas. Entomol.* 19: 269-273.
- Pascual-Villalobos MJ 1996. Evaluation of the insecticidal activity of *Chrysanthemum coronarium* L. plant extracts. *Bol San Veg Plagas* 22: 411-420.
- Perez SG, Ramos-Lopez MA, Zavala-Sanchez MA, Cardenas-Ortega NC 2010. Activity of essential oils as a biorational alternative to control coleopteran insects in stored grains. *J. Medicinal Plants Res.* 4: 2827-2835.
- Rattan RS 2010. Mechanism of action of insecticidal secondary metabolites of plant origin. *Crop Protect.* 29: 913-920.
- Regnault-Roger C, Hamraoui A 1995. Fumigant toxic activity and reproductive inhibition induced by monoterpenes on *Acanthoscelides obtectus* (Say) (Coleoptera), a bruchid of kidney bean (*Phaseolus vulgaris* L.). *J. Stored Prod. Res.* 31: 291-299.
- Rozman V, Kalinovic I, Korunic Z 2007. Toxicity of naturally occurring compounds of Lamiaceae and Lauraceae to three stored-product insects. *J. Stored Prod. Res.* 43: 349-355.
- Sertkaya E, Kaya K, Soylu S 2010. Chemical compositions and insecticidal activities of the essential oils from several medicinal plants against the cotton whitefly, *Bemisia tabaci*. *Asian J. Chem.* 22: 2982-2990.
- Sertkaya E 2013. Fumigant toxicity of the essential oils from medicinal plants against bean weevil, *Acanthoscelides obtectus* (Say) (Coleoptera: Bruchidae). *Asian J. Chem.* 25: 553-555.
- Shukla RS, Singh P, Prakash B, Kumar A, Mishra PK, Dubey NK 2011. Efficacy of essential oils of *Lippia alba* (Mill.) N.E. Brown and *Callistemon lanceolatus* (Sm.) Sweet and their major constituents on mortality, oviposition and feeding behaviour of pulse beetle, *Callosobruchus chinensis* L. *J. Sci Food Agric* 91: 2277-2283.
- Soylu EM, Soylu S, Kurt S 2006. Antimicrobial activities of the essential oils of various plants against tomato late blight disease agent *Phytophthora infestans*. *Mycopathologia* 161: 119-128.
- Soylu S, Yigitbas H, Soylu EM, Kurt S 2007. Antifungal effects of essential oils from oregano and fennel on *Sclerotinia sclerotiorum*. *J. Appl. Microbiol.* 103: 1021-1030.
- Soylu S, Evrendilek GA, Soylu EM 2009. Chemical compositions and antibacterial activities of bitter fennel (*Foeniculum vulgare* Mill. var. vulgare) and dill (*Anethum graveolens* L.) essential oils against the growth of food-borne and seed-borne plant pathogenic bacteria. *Italian J. Food Sci.* 21: 347-355.
- Soylu EM, Kurt S, Soylu S 2010. *In vitro* and *in vivo* antifungal activities of the essential oils of various plants against tomato grey mould disease agent *Botrytis cinerea*. *Int. J. Food Microbiol.* 143:183-189.
- Suthisut D, Fields PG, Chandrapatya A 2011. Contact toxicity, feeding reduction, and repellency of essential oils from three plants from the ginger family (Zingiberaceae) and their major components against *Sitophilus zeamais* and *Tribolium castaneum*. *J. Econ. Entomol.* 104: 1445-1454.
- Toudert-Taleb K, Hedjal-Chebheb M, Hami H, Debras JF, Kellouche A 2014. Composition of essential oils extracted from six aromatic plants of Kabyle origin (Algeria) and evaluation of their bioactivity on *Callosobruchus maculatus* (Fabricius, 1775) (Coleoptera: Bruchidae). *African Entomol.* 22: 417-427.
- Tunç İ, Erler, F 2003. Repellency and repellent stability of essential oil constituents against *Tribolium confusum*. *Z. Pflanzenk. Pflanzen.* (Journal of Plant Diseases and Protection), 110: 394-400